

Narrative Review

Dry needling: a literature review with implications for clinical practice guidelines¹

James Dunning^{1,2}, Raymond Butts^{3,4}, Firas Mourad⁵, Ian Young⁶, Sean Flanagan⁷, Thomas Perreault⁸

¹Nova Southeastern University, Ft. Lauderdale, FL, USA, ²Alabama Physical Therapy & Acupuncture, Montgomery, AL, USA, ³University of South Carolina, Columbia, SC, USA, ⁴Palmetto Health Research Physical Therapy Specialists, Columbia, SC, USA, ⁵Sportlife Physiotherapy, Montichiari, Italy, ⁶Spine & Sport, Savannah, GA, USA, ⁷OneAccord Physical Therapy, Casa Grande, AZ, USA, ⁸Portsmouth-Newington Physical Therapy, Portsmouth, NH

Background: Wet needling uses hollow-bore needles to deliver corticosteroids, anesthetics, sclerosants, botulinum toxins, or other agents. In contrast, dry needling requires the insertion of thin monofilament needles, as used in the practice of acupuncture, without the use of injectate into muscles, ligaments, tendons, subcutaneous fascia, and scar tissue. Dry needles may also be inserted in the vicinity of peripheral nerves and/or neurovascular bundles in order to manage a variety of neuromusculoskeletal pain syndromes. Nevertheless, some position statements by several US State Boards of Physical Therapy have narrowly defined dry needling as an 'intramuscular' procedure involving the isolated treatment of 'myofascial trigger points' (MTrPs).

Objectives: To operationalize an appropriate definition for dry needling based on the existing literature and to further investigate the optimal frequency, duration, and intensity of dry needling for both spinal and extremity neuromusculoskeletal conditions.

Major findings: According to recent findings in the literature, the needle tip touches, taps, or pricks tiny nerve endings or neural tissue (i.e. 'sensitive loci' or 'nociceptors') when it is inserted into a MTrP. To date, there is a paucity of high-quality evidence to underpin the use of direct dry needling into MTrPs for the purpose of short and long-term pain and disability reduction in patients with musculoskeletal pain syndromes. Furthermore, there is a lack of robust evidence validating the clinical diagnostic criteria for trigger point identification or diagnosis. High-quality studies have also demonstrated that manual examination for the identification and localization of a trigger point is neither valid nor reliable between-examiners.

Conclusions: Several studies have demonstrated immediate or short-term improvements in pain and/or disability by targeting trigger points (TrPs) using in-and-out techniques such as 'pistoning' or 'sparrow pecking'; however, to date, no high-quality, long-term trials supporting in-and-out needling techniques at exclusively muscular TrPs exist, and the practice should therefore be questioned. The insertion of dry needles into asymptomatic body areas proximal and/or distal to the primary source of pain is supported by the myofascial pain syndrome literature. Physical therapists should not ignore the findings of the Western or biomedical 'acupuncture' literature that have used the very same 'dry needles' to treat patients with a variety of neuromusculoskeletal conditions in numerous, large scale randomized controlled trials. Although the optimal frequency, duration, and intensity of dry needling has yet to be determined for many neuromusculoskeletal conditions, the vast majority of dry needling randomized controlled trials have manually stimulated the needles and left them *in situ* for between 10 and 30 minute durations. Position statements and clinical practice guidelines for dry needling should be based on the best available literature, not a single paradigm or school of thought; therefore, physical therapy associations and state boards of physical therapy should consider broadening the definition of dry needling to encompass the stimulation of neural, muscular, and connective tissues, not just 'TrPs'.

Keywords: Dry needling, Literature review, Trigger point, Practice guidelines, Physical therapy

Introduction

Local injection therapies, often referred to as 'wet needling', use hollow-bore needles to deliver corticosteroids, anesthetics, sclerosants, botulinum toxins, or other agents.^{1,2} In contrast, 'dry needling' refers to the

¹This paper should be attributed to: College of Health Care Sciences, Nova Southeastern University, Ft. Lauderdale, FL, USA.

Correspondence to: James Dunning, 1036 Old Breckenridge Lane, Montgomery, AL 36117, USA. Email: jamesdunning@hotmail.com

insertion of thin monofilament needles, as used in the practice of acupuncture, without the use of injectate.³⁻⁶ Dry needling is typically used to treat muscles, ligaments, tendons, subcutaneous fascia, scar tissue, peripheral nerves, and neurovascular bundles for the management of a variety of neuromusculoskeletal pain syndromes.^{3,6,7} Given the broad base of international literature presently available on the technique, it is particularly concerning that the primary US-based, National Physical Therapy Association⁸ and several State Boards of Physical Therapy⁹⁻¹³ have recently narrowed their definition of dry needling to an ‘intramuscular’ procedure, i.e. the insertion of needles into nodules within taut bands of muscle, more commonly referred to as ‘trigger points’ (TrPs) or ‘myofascial trigger points’ (MTrPs). More specifically, these professional organizations have equated the procedure of dry needling with the term ‘intramuscular manual therapy’ (IMT) or ‘trigger point dry needling’ (TDN). Certainly, IMT, or the insertion of needles into TrPs within muscle bellies, is one aspect of dry needling; however, IMT or TDN should not be used synonymously with the term dry needling. In addition to TrPs within muscular tissue, an extensive body of literature, including both peer-reviewed articles and randomized controlled trials, supports the insertion of dry needles throughout the body at non-TrP sites for the purpose of reducing pain and disability in patients with neuromusculoskeletal conditions. Just as the Maitland,¹⁴ Kaltenborn,¹⁵ and McKenzie¹⁶ approaches are each unique brands of manual therapy, ‘TDN’ and ‘IMT’ are individual aspects of dry needling. Each describes a single framework, paradigm, or approach that falls under the much larger field of dry needling.^{5,6,9,10,17-25}

Dry Needling Targets Neural, Muscular, and Connective Tissues, Not Just TrPs

The first, peer-reviewed journal article on dry needling published by a Western, medical physician⁶ did not limit needle insertion to muscular ‘TrPs’; rather, the 241-patient study reported that only 2 of the 14 target structures were muscular TrPs.⁶ The other structures needled included ligaments, scar tissue, tendons, bones, and teno-osseous insertion sites, all of which are types of connective tissue.⁶ In addition, ‘a high density of neurovascular structures’ has been found at dry needling target sites.²⁶

As early as 1977, Melzack *et al.*²⁷ stated that ‘Trigger points are firmly anchored in the anatomy of the neural and muscular systems... and the stimulation of particular nerves or tissues by needles could bring about an increased input to the central biasing mechanism, which would close the gates to [pain] inputs from selected body areas’.²⁷ In a more recent commentary titled ‘Treatment of Myofascial Pain

Syndrome’, Hong stated that the purpose of ‘the fast-in and fast-out needle technique’^{26,28-30} in a fan or cone shape is to ‘ensure that all or most sensitive loci (i.e. tiny nerve endings) are encountered’.²⁸ Widely considered one of the world authorities on both dry needling practice and experimental research, Hong further reported that ‘high-pressure stimulation by needling’^{26,28,31-33} or ‘mechanical irritation of the needle’^{26,34} to ‘multiple sensitive loci or nociceptors within the same MTrP’^{26,28,31-34} likely ‘elicits a local twitch response that subsequently provides a very strong neural impulse to the MTrP circuit to break the vicious cycle so that MTrP pain is relieved’.^{26,28,33,34} Interestingly, ‘bilateral or mirror image evoked potentials’ or ‘local twitch responses’³⁴ (LTRs) have been recorded in the contralateral trapezius during unilateral dry needling stimulation of active TrPs on the ipsilateral, symptomatic trapezius.³⁴ Audette *et al.*³⁴ concluded, ‘the LTR is generated by inducing a spinal reflex that involves sensory input to the spinal cord by mechanical irritation of the needle at sensitive loci in the TrP which then results in a motor, efferent response of the alpha motor neuron pool’.³⁴ The presence of bilateral LTRs during unilateral dry needling ‘argues strongly for a central abnormality rather than a purely peripheral [muscular] abnormality in patients with active MTrPs’.³⁴

Thus, within the context of myofascial pain syndromes, dry needles likely come in contact with both muscle and connective tissue.⁶ It is held by several research teams that needles inserted into MTrPs likely touch, tap, or prick tiny nerve endings or neural tissue for the purpose of pain reduction. Based on the most recent literature,^{6,21,26,28-37} the definition of dry needling by the American Physical Therapy Association (APTA) and State Boards of Physical Therapy should therefore clearly articulate that dry needling encompasses stimulation of neural,^{26,28-31,33,36,38-54} muscular,^{5,17,18,21,26,29,30,33,48,55-69} and connective tissues,^{6,48-54,70-72} and not just ‘TrPs’.^{73,74}

2013 APTA Definition: Dry Needling Targets Both Muscular and Connective Tissues

Page two of the January 2012 Resource Paper⁸ titled, *Physical Therapists & the Performance of Dry Needling* defined dry needling as ‘an invasive technique used by physical therapists (where allowed by state law) to treat myofascial pain that uses a dry needle, without medication or injection, which is inserted into areas of the muscle known as trigger points...’⁸ Notably, an updated February 2013 APTA version of this document titled, *Description of Dry Needling in Clinical Practice: an Educational Resource Paper*, defines dry needling as including both muscular and connective tissues, but not neural stimulation.⁷⁵ The paper reads, ‘dry needling is a

skilled intervention that uses a thin filiform needle to penetrate the skin and stimulate underlying myofascial trigger points, muscular, and connective tissues for the management of neuromusculoskeletal pain and movement impairments'.⁷⁵ Interestingly, the 2013 definition implies that only muscle and connective tissue may be targeted by dry needles in the treatment of neuromusculoskeletal pain.^{26,28,31-34} Yet, nerve endings, referred to as 'sensitive loci' or 'neurovascular bundles' are clearly supported as needle stimulation targets in the existing literature.²⁶ How can physical therapists treat the 'neuro' in neuromusculoskeletal with dry needles without placing needles near, beside, or onto neural tissue (i.e. peri-neural needling)? By adding neural targets to the position statement, the APTA would provide a stronger and more evidence-based framework for state boards to follow.

Narrowly Focused Position Statements by State Boards of Physical Therapy

To date, a number of State Physical Therapy Boards⁹⁻¹³ continue to have limited definitions of dry needling. As a result, many physical therapists believe that they are only able to treat patients by targeting intramuscular TrPs with dry needles, as their respective State Boards do not authorize them to insert needles into connective tissues (e.g. ligaments, tendons, teno-osseous junctions, musculotendinous junctions, scar tissue) or to perform peri-neural or peri-neurovascular needling. Several State Boards of Physical Therapy have issued narrowly focused position statements on dry needling:

- *Mississippi (Effective: 10 September 2012)*: The Mississippi State Board of Physical Therapy updated its regulations to read, 'Intramuscular manual therapy is a physical intervention that uses a filiform needle no larger than a 25-gauge needle to stimulate trigger points, diagnose and treat neuromuscular pain and functional movement deficits; is based upon Western medical concepts; requires an examination and diagnosis, and treats specific anatomic entities selected according to physical signs'.¹⁰
- *North Carolina (Effective: 14 June 2012)*: The North Carolina Board of Physical Therapy Examiners revised its definition of dry needling to read, 'Intramuscular Manual Therapy (IMT), which is generally referred to as dry needling, is defined as a technique to treat myofascial pain using a dry needle (without medication) that is inserted into a trigger point with the goal of releasing/inactivating the trigger points and relieving pain'.⁹
- *Louisiana (Effective: 20 October 2011)*: Rule 311 of the Louisiana Physical Therapy State Board uses the terms 'dry needling' and 'intramuscular dry needling' interchangeably, and as a result, requires a physical therapist to 'successfully complete...no fewer than 50 hours of face-to-face instruction in intramuscular dry needling treatment and safety'.¹¹
- *Nebraska (Effective: 20 June 2011)*: The Nebraska State Board of Physical Therapy opined, 'Dry needling is a mechanical modality technique used to treat myofascial pain that uses a dry needle, without medication, that is inserted into a trigger point with the goal of releasing/inactivating the trigger points'.¹²
- *Colorado (Effective: 30 June 2012)*: The Colorado State Physical Therapy Board opined, 'Dry needling (also known as Trigger Point Dry Needling) is a physical intervention that uses a filiform needle to stimulate trigger points, diagnose and treat neuromuscular pain and functional movement deficits; is based upon Western medical concepts; requires an examination and diagnosis, and treats specific anatomic entities selected according to physical signs. Dry needling does not include the stimulation of auricular or distal points'.¹³

Whether these State Boards of Physical Therapy⁹⁻¹³ will update their position statements and/or practice acts to include both neural and connective tissue as possible target structures for dry needling remains to be seen. However, Arizona has recently provided a reason to be optimistic. In January 2013, the Arizona Physical Therapy Association set the standard by becoming the first US-based, physical therapy association to explicitly recognize neural, muscular, and connective tissue as target sites for dry needling.⁷⁶ The definition of dry needling that appears in the Arizona position statement reads,

*It is the position of the Arizona Physical Therapy Association that dry needling for the management of neuromusculoskeletal conditions is consistent with the scope of practice of licensed physical therapists in Arizona. Dry Needling is a skilled intervention performed by a physical therapist that uses a thin filiform needle to penetrate the skin and stimulate underlying neural, muscular and connective tissues for the evaluation and management of neuromusculoskeletal pain and movement impairments.*⁷⁶

Page two of the January 2012 APTA Resource Paper⁸ titled, 'Physical Therapists & the Performance of Dry Needling', also states, 'Preliminary research supports that dry needling improves pain control, reduces muscle tension, normalizes biochemical and electrical dysfunction of motor endplates, and facilitates an accelerated return to active rehabilitation'. However, although the reference list of the 141-page resource paper⁸ cited four systematic reviews,^{17,19,24,56} one clinical review,²¹ and one unpublished evidence summary,⁷⁷ the 23 randomized controlled trials that were reportedly reviewed and rated by a single expert on a 0-5 scale for quality and level of support for dry needling, were not individually identified or specifically referenced in any way.⁸ Moreover, while the 'median quality of the research was 4' and the 'median support for dry needling was 3', the specific studies included in the analysis remains a mystery.⁸ This resource paper⁸ also failed to describe available

literature on neuromusculoskeletal conditions that supports the use of ‘TrP’ dry needling.

It is concerning that the APTA^{8,75} and several State Boards of Physical Therapy^{9–13} continue to omit neural and/or connective tissue as possible target structures when defining dry needling in official position statements. Perhaps position statements that compartmentalize dry needling to ‘IMT’ – i.e. confine it to muscle bellies, tight knots, or ‘myofascial TrPs’,^{8–11} – fit better with the traditional physical therapy framework. Given that other professions continue to question whether dry needling is within the scope of practice of a physical therapist, declaratory statements by State Boards of Physical Therapy that operationalize dry needling with an ‘intramuscular’^{9–13} framework may be a more defensible position. Certainly, ‘nerve points’^{42,43} sound a little ‘Eastern’ to some, and may, at face value, be confused with traditional Chinese acupuncture; however, dry needling neither attempts to move *qi* along meridians, nor does it rely on diagnoses from traditional Chinese acupuncture or Oriental medicine.^{78,79}

Trigger Point Needling is Only One Kind of Dry Needling

In 2009, the APTA recommended ‘IMT’ as the term to be used by physical therapists to describe the intervention of dry needling.⁸ The following excerpt is from page two of the January 2012 Resource Paper titled, ‘Physical Therapists & the Performance of Dry Needling’:

Dry needling is an invasive technique used by physical therapists (where allowed by state law) to treat myofascial pain that uses a dry needle, without medication or injection, which is inserted into areas of the muscle known as TrPs. A TrP describes a taught band of skeletal muscle located within a larger muscle group. Trigger points can be tender to the touch and can refer pain to distant parts of the body. Physical therapists utilize dry needling with the goal of releasing/inactivating the TrPs and relieving pain.⁸

This definition seems to suggest that physical therapists should only insert needles into muscle bellies in an attempt to inactivate TrPs. This recommendation makes sense considering Itoh *et al.*⁸⁰ found that ‘TrP acupuncture’ of the splenius capitis, upper trapezius, levator scapulae, sternocleidomastoid, suboccipitals, scaleni and paravertebrals was more effective in the short-term (i.e. at week 13) at reducing pain and disability than classical or sham acupuncture in patients with chronic, non-radiculopathic neck pain. In contrast, Tough *et al.* concluded in a 2009 systematic review and meta-analysis that ‘there is limited evidence deriving that deep needling directly into myofascial trigger points has an overall treatment effect when compared to standardized care’.²⁴

Moreover, ‘the result of the meta-analysis of direct dry needling [for myofascial trigger point pain] compared to placebo control treatment did not attain statistical significance’.²⁴ There is a paucity of high-quality evidence to underpin the use of direct dry needling into myofascial TrPs for the purpose of short-/long-term pain and disability reduction in patients with musculoskeletal pain syndromes.^{18,20,23,24,80} Rather, the vast majority of the dry needling literature for the management of neuromusculoskeletal conditions has demonstrated that targeting TrPs (i.e. muscle tissue) and non-TrP structures (i.e. neural and connective tissues) is as effective as needling only non-TrP structures in reducing pain and disability in patients with the following conditions: knee osteoarthritis,^{50–52,81–90} hip osteoarthritis,^{53,91–94} piriformis syndrome,⁶⁰ carpal tunnel syndrome,^{36,37,40,41,44,45,47} migraine,^{95–101} tension type headache,^{97,98,102} temporomandibular disorder,^{59,67,94,103–107} shoulder pain,^{108–110} neck pain,^{23,24,29,35,61,63–65,80,94,111–114} low back pain,^{4,5,19,20,66,69,94,115–125} and plantar fasciitis.^{48,49,54,56,126,127}

While many of these studies directly use the term ‘dry needling’,^{5,18,21,26,33,56,64,67,104,110} in their methodology, other studies use alternative terminology such as ‘deep dry needling’,⁶⁷ ‘superficial dry needling’,^{55,128} ‘TrP acupuncture’,^{20,80} ‘paraspinal needling’,⁶⁴ ‘intramuscular and nerve root needling’,⁶⁵ ‘needle electrical intramuscular stimulation’,¹²⁹ ‘needle release’,¹¹³ ‘acupuncture needling’,^{63,113} ‘needling therapy’,¹⁷ ‘acupuncture’,^{36–38,40,41,47,49–55,61,69,72,81–88,90,92,94–100,105–108,112,115–117,119–123,130–135} and ‘electroacupuncture’.^{37,46,93,126,127,136–139} Some studies have even chosen to use both ‘acupuncture and dry needling’^{19,23,24,62,77} in their titles. Importantly, none of the studies used medicine and/or injectate in conjunction with their needling procedure; therefore, all studies fit within the strict definition of dry needling, regardless of terminology. To our knowledge, no published randomized controlled trials, to date, have used the words ‘IMT’ to describe a dry needling intervention for any condition.

Interestingly, the most common term used to describe dry needling is ‘acupuncture’. Physiotherapists and/or medical physicians^{114–116,119,140–144} within both government administered national health services and mainstream university health systems^{53,90,97–99,114,123,133,134} in the UK,^{49,51,52,84,100,101,131,132} Canada,¹¹⁴ USA and Germany^{53,90,97–99,123,133,134} use the term ‘acupuncture’ to describe dry needling methodologies. The same is true of articles published in mainstream, highly respected journals, including the *British Medical Journal*,^{84,89,100,101,112,124} *European Journal of Pain*,^{24,131} *Archives of Physical Medicine & Rehabilitation*,^{17,22,60,109,145,146} *Pain*,^{6,44,87,111,123,125,133,136,147} *Annals of Internal Medicine*,^{4,81,86,88,117,132} *Headache*,^{95,96} and *Rheumatology*.^{35,51,82,85}

Spine,^{5,19,66,114,118,119,121,135} and *Cochrane Database of Systematic Reviews*.^{62,108} Even the *Physical Therapy Journal* sponsored by the APTA has used acupuncture and dry needling interchangeably in a recent publication.⁵² It would therefore be a mistake to ignore the findings of high-quality, randomized controlled trials,^{37,40,53,54,63,81,84,89,90,92,99,100,107,114,115,119,125,132,148} systematic reviews,^{17,19,23,56,83,97,121,149} meta-analyses,^{24,85} Cochrane reviews,^{62,108,150} the British practice guidelines,¹²⁴ the European practice guidelines,^{151,152} and the joint clinical practice guidelines from the American College of Physicians and the American Pain Society¹¹⁷ simply because they used the term ‘acupuncture’ instead of dry needling in their title and/or methods section. Moreover, ignoring RCTs published by PhD and licensed acupuncturists (LAc) in well-respected, peer-reviewed journals would be shortsighted. By disregarding these studies, physical therapists may limit their ability to support the use of dry needling in clinical practice for treating knee osteoarthritis, shoulder pain, carpal tunnel syndrome, migraine headache, tension type headache, hip pain, low back pain, and neck pain.

Physical Therapists Should Not Ignore the Findings of Western Acupuncture Trials

‘Acupuncture’ literally translates to ‘*needle penetration*’ (i.e. ‘acu’=needle; ‘puncture’=penetration), and the vast majority of the ‘acupuncture’ trials are not claiming to move *qi* along meridians or channels. Furthermore, the vast majority of the so-called ‘acupuncture’ RCTs have used Western medical diagnoses (not traditional Chinese or Oriental medicine^{78,79} diagnoses such as *bi* syndrome, *qi*, blood stagnation, and kidney *yang* deficiency^{148,153}) such as chronic neck pain,^{23,24,29,35,61,63–65,80,94,111–114} plantar fasciitis,^{48,49,54,56,126,127} knee osteoarthritis,^{50–52,81–90} and carpal tunnel syndrome.^{36,37,40,41,44,45,47} These same ‘acupuncture’ trials have investigated the efficacy of inserting thin filiform needles (without medicine) into ‘Ah-Shi points’ (in Chinese ‘auwh’ means ‘that’s where it hurts’ or ‘that’s it’; synonymous with trigger points^{57,58,154}) and/or non-trigger point locations. Thus, while the terminology, theoretical constructs, and philosophies are different, the actual procedure of inserting thin monofilament needles, as used in the practice of acupuncture, without the use of injectate is very similar across professions.³ The use of McKenzie and May¹⁶ exercises by a chiropractor does not make him or her a physical therapist; likewise, the use of dry needles by a physical therapist does not make him or her an acupuncturist. Rather, these techniques are shared procedures among a variety of healthcare professions. As an additional comparison, physical therapist researchers do not ignore or exclude studies published by MDs, DOs, and DCs when citing

references to support the use of spinal manipulation treatments for a variety of neuromusculoskeletal conditions simply because the authors consider the techniques as ‘chiropractic’ or ‘osteopathic manipulations’. Physical therapists should therefore not ignore the findings of large scale randomized controlled trials available in the Western or biomedical ‘acupuncture’ literature that use the same ‘dry needles’ to treat patients with neuromuscular conditions.^{49,51–53,84,90,97–101,114,123,131–134}

Poor Inter-Examiner Reliability for TrP Location

Barbero *et al.*¹⁵⁵ reported ‘moderate to high’ intra-rater reliability of an experienced physiotherapist for determining the location of MTrPs within the upper trapezius muscle. However, in a recent systematic review on the reliability of physical examination for the diagnosis of myofascial TrPs, Lucas *et al.*¹⁴⁹ concluded, ‘There is no accepted reference standard for the diagnosis of trigger points, and data on the reliability of physical examination for trigger points is conflicting’. In addition, a predictable pattern of pain referral and the local twitch response are each no longer considered to be sufficient or necessary for the diagnosis of a TrP.^{22,25,149} After reviewing nine studies on reliability, Lucas *et al.*¹⁴⁹ further concluded:

*None of the nine studies in this systematic literature review specifically reported inter-rater reliability estimates for the identification of the location of active trigger points in symptomatic participants.... At present, there is no data on the reliability of pinpointing the exact location of active trigger points.... The existing data on reliability pertain only to agreeing if a muscle has the signs of a trigger point and not the exact location of the taut band or the nodule within the taut band.*¹⁴⁹

Lew *et al.*¹⁵⁶ reported that the inter-examiner agreement was only 21%, and Myburgh *et al.*¹⁵⁷ found error rates of 3.3–6.6 cm among examiners attempting to identify the specific location of TrPs in the upper trapezius muscle. In another recent literature review, Myburgh *et al.*²² found poor inter-examiner reliability of manual palpation of TrPs in various muscle groups. Only ‘tenderness’ of the upper trapezius, not the actual location of the TrP, was found to be moderately reliable. In 2011, Myburgh *et al.*¹⁵⁷ similarly reported ‘good agreement between experienced practitioners’ for the ‘presence or absence’ of a clinically relevant TrP in the upper trapezius muscle. However, this study, like the many others,^{158–162} failed to investigate the inter-rater reliability for determining the specific location of the TrP within the target muscle. If clinicians are not able to reliably identify TrPs, they may not be able to consistently activate/deactivate them by penetrating the nodule within the taut band, as is commonly

taught by Travell and Simons.^{73,74} In fact, in a recent systematic review, Tough *et al.*²⁵ concluded, ‘There is a lack of robust empirical evidence validating the clinical diagnostic criteria [for TrP identification or diagnosis] proposed by both Travell and Simons (1999) and Fischer (1997)’.

High-quality evidence suggests that manual examination for the identification of the specific location of the ‘TrP’ is not a valid^{22,25,157} or reliable^{22,147,149,156} process between-examiners. Poor reliability of TrP identification also makes it rather challenging, if not impossible, for Physical Therapists to be in compliance with the APTA⁸ and several State Physical Therapy Boards.^{93,120,127,129,155} As Lucas *et al.*¹⁴⁹ summarizes, ‘It is not yet evident that examiners can agree on the precise location of an active TrP; hence, they cannot be relied upon to accurately insert the needle into the nodule of the taut band’. Perhaps ‘TrP dry needling’, or ‘IMT’,^{8–13} as performed by physical therapists, does not meet the criteria of evidence-based practice (i.e. a valid and reliable diagnostic and therapeutic intervention).

Ah-Shi Acupuncture Points and TrPs: Different Theories, Similar Locations

Although ‘IMT’ or ‘TrP needling’ involves insertion of needles into muscle bellies, Melzack *et al.*²⁷ reported:

*a high degree (71%) of correspondence between MTrPs and acupuncture points, and it is very likely that all MTrPs are Ah-Shi acupuncture points... [however], trigger points are firmly anchored in the anatomy of the neural and muscular systems, while acupuncture points are associated with an ancient conceptual but anatomically non-existent system of meridians, which carry Yin (spirits) and Yang (blood).*²⁷

Melzack *et al.*²⁷ further concluded, ‘trigger points and acupuncture points for pain, though discovered independently, and labeled differently, represent the same phenomenon and can be explained in terms of the same underlying neural mechanisms’. More recently, Dorsher and Fleckenstein,⁵⁷ both medical physicians, found that 238 (93.3%) of 255 common MTrPs anatomically corresponded with classical acupoints. Furthermore, ‘the marked correspondences of the pain indications (up to 97%) and somatovisceral indications (up to 93%) of anatomically corresponding common MTrP-classical acupoint pairs provide a second, clinical line of evidence that trigger points and acupuncture points likely describe the same physiologic phenomena’.⁵⁸ Dorsher further demonstrated a strong correspondence between the distributions of the myofascial referred-pain patterns^{73,74} and acupuncture meridians^{78,79} of anatomically corresponding common MTrP-classical acupoint pairs.

Biomechanical, Chemical, Endocrinological, and Vascular Effects of Dry Needling

The biomechanical,^{70–72,163} chemical,^{146,164} and vascular^{129,138,140,165,166} effects of needling either superficial⁵⁵ subcutaneous tissue (non-muscular) or deep^{23,24,73,74} intramuscular tissue without injectate have been well documented. Improved microcirculation around the knee joint has been demonstrated following ‘dry’ needling into non-TrP locations,¹³⁸ and improved muscle blood flow has been found following ‘manual acupuncture’ in the lower extremities.^{167,168} Following needle electrical intramuscular stimulation targeting MTrPs in patients with shoulder and cervical myofascial pain syndrome, microcirculation above the area of the MTrPs was found to have increased from 127.3 to 310.2 BPU per Laser Doppler Flowmetry, a greater than two-fold increase. Furthermore, low blood flow seemed to correlate with pain intensity,¹²⁹ suggesting that reduced microcirculation plays a role in the pathophysiology of myofascial pain syndrome.^{166,169} Electrical dry needling in patients with knee osteoarthritis has also resulted in endocrinological changes, including increases in beta-endorphins and decreases in cortisol.¹³⁶

Non-Trigger Point Dry Needling in Knee Osteoarthritis and Carpal Tunnel Syndrome: Highly Effective

There is overwhelming evidence from randomized controlled trials, systematic reviews, and meta-analyses that dry needling the knee joint without targeting specific TrPs is effective at reducing pain and disability in patients with chronic knee osteoarthritis.^{50–52,81–85,89,90,131} Likewise, there is robust evidence that peri-neural needling of non-TrP structures helps reduce pain and disability while improving sensory and motor nerve conduction velocities. Peri-neural needling has also been shown to stimulate microcirculation in patients with mild to moderate carpal tunnel syndrome.^{36,37,40,41,47} Such findings suggest that organizations that teach physical therapists a TrP ‘search and destroy’ method of dry needling may not be evidence based. Rather, therapists may be much more effective treating conditions such as osteoarthritis and carpal tunnel syndrome by focusing on non-TrP locations such as neural, connective, and muscle tissue.

Dry Needling for Tendinopathy, Not Just TrPs

Several recent studies^{7,163,165,168,170–172} support the use of dry needling in the management of tendinopathy; more specifically, dry needling has been shown to positively influence tendon healing by increasing blood flow via local vasodilation^{165,167,168,170} and collagen proliferation.^{163,171} It is theorized that the lack of blood flow and subsequent hypoxic environment contributes to ‘tendon dysrepair’¹⁷³ and leads to the

pain and disability associated with chronic tendinopathy.⁷ In comparison to a superficial heating modality, and following dry needling using ‘vertical pecking’ between the tendon and its sheath, Kubo *et al.*¹⁶⁵ found statistically significant increases in blood flow and oxygen saturation levels around the Achilles tendon. Following dry needling to burn-injured mice, and when compared with conventional dressing, Lee *et al.*¹⁷¹ found significantly greater basic fibroblastic growth factor and accelerated tissue healing rates. Likewise, Langevin *et al.*¹⁶³ reported significantly greater fibroblastic activity following rotational needle manipulation in a mouse model. Clearly, dry needling both superficial and deep non-TrP locations results in significant mechanical,^{70–72,163} chemical,^{146,164} endocrinological,¹³⁶ microvascular,^{138,165,167,168,171} neural,^{36,41,47} and central effects^{38,39,44,46,130,137,140} (i.e. activation of the descending pain inhibitory systems, cortex, hypothalamus, and inactivation of the limbic system per recent fMRI, and PET studies). As such, the physical therapy profession must re-evaluate whether it is truly ‘best practice’ to limit dry needling to muscle TrPs.

There is Limited Evidence Supporting the Effectiveness of Deep TDN

Compared to sham or placebo treatments, a recent systematic review and meta-analysis¹⁷⁴ recommended ‘dry needling [directly into MTrPs] for decreasing pain in patients with upper quarter myofascial pain syndrome’; however, the data in this meta-analysis was based on three ‘immediately after’ and two ‘at 4-weeks’ post-treatment studies. More specifically, although the findings of Kietrys *et al.*¹⁷⁴ support the immediate and short-term effectiveness of dry needling, it does not provide any evidence for the long-term effects of direct TDN on pain or disability in patients with upper quarter myofascial pain syndrome. There are three trials^{20,80,113} that seem to be frequently cited by clinicians in support of deep TDN; however, all three studies have significant methodologic limitations. First, Ma *et al.*¹¹³ used a ‘mini-scalpel needle release’ surgical procedure instead of monofilament dry needling in an effort to remove or disrupt a TrP within the upper trapezius. As such, the study¹¹³ should not be cited in support of ‘sparrow pecking’, ‘pistoning’, ‘fast-in fast-out’

intramuscular and deep TDN in patients with myofascial pain syndrome. Second, although Itoh *et al.*²⁰ found that deep (20 mm) TrP acupuncture of myofascial TrPs is more effective than either superficial (3 mm) TrP acupuncture or standard acupuncture, the conclusions should be cautiously considered given the following methodological shortcomings: (1) the sample contained just 35 patients that were further split into three smaller groups, (2) the longest outcome measures for pain and disability were taken just 4 weeks after the final needling session, (3) the improvements in pain seen in the deep TrP acupuncture group was ‘reversed’ by the end of the 3-week no treatment interval, (4) the between-group differences for pain and disability were not statistically significantly different by week 13. Third, after comparing TrP, standard, and sham acupuncture, Itoh *et al.*⁸⁰ concluded that TrP acupuncture is more effective than standard (traditional/classical) acupuncture and non-penetrating sham acupuncture for reducing pain and disability in patients with chronic, non-radiculopathic neck pain. However, the 40-patient sample size was divided among three smaller groups, and outcome measures for pain and disability were taken only 3 weeks post-treatment.

Multiple Needles Should Be Left *In Situ* for 10–30 minutes

Although the optimum dosage^{50,51,83,85,124} (frequency of treatment sessions per week or month), duration (length of time the needles should remain *in situ*), and intensity (the number of needles used and degree of manual manipulation or electrical stimulation) has yet to be determined^{23,24,50,51,83,85,114,135,141} for many neuromusculoskeletal conditions, the vast majority of ‘dry’ needling randomized controlled trials attempt to elicit a *deqi* response.^{142,143} *Deqi* has been defined as a dull ache, heaviness, distension, numbness, tingling, cramping, pressure, fullness, spreading, warmth, or coolness.^{142,143} In addition, most trials have left multiple needles *in situ* for between 5 and 40-minute durations; moreover, a recent systematic review and meta-analysis¹⁹ within the framework of the Cochrane collaboration concluded, ‘for low back pain of any duration, needle retention for about 10 minutes is better than removal immediately after insertion’.¹⁹ Table 1 provides the durations and

Table 1 Duration needles left *in situ* for neuromusculoskeletal condition.

Neuromusculoskeletal condition	Duration needles left <i>in situ</i>
Low back pain	10 minutes, ²⁰ 15 minutes, ⁶⁸ 20 minutes ^{116,119,125,144} or 30 minutes. ^{69,123,139}
Carpal tunnel syndrome	30 minutes, ^{37,40,41} 40 minutes ⁴⁷ or 60 minutes. ³⁶
Plantar fasciitis	5 minutes, ^{56,186,187} 15 minutes ⁴⁹ or 20 minutes. ^{126,127}
Knee osteoarthritis	20 minutes ^{51,81,82,87,88,136} or 30 minutes. ^{88–90,138}
Shoulder pain	5 minutes ¹⁸ or 10 minutes. ¹¹⁰
Neck pain	10 minutes, ⁸⁰ 20 minutes ^{61,132,141,188} or 30 minutes. ^{111,112,133,189}
Headaches	20 minutes ⁹⁵ or 30 minutes. ^{96,98–101}

specific studies that multiple needles were left *in situ* for a variety of neuromusculoskeletal conditions.

Even when ‘pistoning’, ‘sparrow pecking’, or ‘fast-in-and-out’ maneuvers were used with the intent of eliciting an LTR in muscular TrPs, multiple needles were often left in place for ten minutes or longer.^{20,80} While several studies^{29,35,59,63,64,67,102,104,175,176} have demonstrated immediate and/or short-term improvements in pain and/or disability following ‘in-an-out’ techniques, to date, there are no high-quality long-term trials^{23,24} to support the practice of immediately removing the needles after pricking TrPs or eliciting LTRs; therefore, this practice should be questioned.

Local, Proximal, and Distal Needling

‘Regional interdependence’ has been defined as ‘the concept that seemingly unrelated impairments in a remote anatomical region may contribute to, or be associated with, the patient’s primary complaint’;¹⁷⁷ that is, the practice of examining and treating muscle, joint or neural impairments that are not local, but are distal or proximal to the patient’s symptoms, has gained widespread acceptance in the orthopedic manual physical therapy community.¹⁷⁷⁻¹⁸³ For example, there are several trials that support the use of thrust manipulation to the cervical, cervicothoracic, thoracic, and upper rib articulations in the treatment of patients with the primary complaint of shoulder pain¹⁷⁸⁻¹⁸³ or shoulder impingement^{184,185}—i.e. treating proximal structures to affect distal symptoms. Likewise, patients with lateral epicondylalgia often seek conservative treatment by a physical therapist. In addition to providing manual therapy to the humero-radial joint and needling the forearm extensor muscles,¹⁷⁶ physical therapists often manipulate the neck and wrist based on supporting evidence in the literature.^{190,191} That is, physical therapists commonly deliver manual therapy both proximal¹⁹⁰ and distal¹⁹¹ to the site of pain. The fact that the Mississippi and Colorado State Boards of Physical Therapy believe that IMT and dry needling, respectively, do not include the stimulation of distal points seems misleading and clinically illogical.¹⁰ Based on this line of reasoning, physical therapists in Mississippi¹⁰ and Colorado¹³ are not allowed to target TrPs in the adductor magnus muscle in patients with ‘generalized internal pelvic pain...or pain shooting up inside the pelvis’⁷⁴ because the pain is located ‘distal’ to the patient’s pain complaint. It is worth noting that this clinical reasoning is directly contrary to that recommended by Travell and Simons. According to Travell and Simons, ‘deep pain in the ipsilateral sacroiliac joint’⁷⁴ may be caused by a TrP in a very ‘distal’ location, namely the soleus muscle. Interestingly, the most recent literature on

myofascial pain syndrome provides clear evidence for needling distal sites, a finding analogous to the acupuncture tradition of treating distal acupoints to influence anatomically remote pain.¹⁵⁴

More recently,³³ ipsilateral or contralateral dry needling to MTrSs (myofascial trigger spots) in the distal gastrocnemius muscle (lower leg) was shown to suppress spontaneous electrical activity (i.e. end plate noise) in MTrSs within the more proximal biceps femoris muscle. Furthermore, after either tibial nerve transection or L5-6 spinal cord transection in rabbits, Hsieh *et al.*³³ found that the remote effects in the bicep femoris disappeared. According to Hsieh *et al.*,³³ the physiologic basis for the remote and/or distal effects of dry needling may be related to activation of the diffuse noxious inhibitory control (DNIC) system, as induced by noxious stimulation provided by the needles. Simply put, ‘the pathway for the remote effect appears to be a spinal reflex’.³³ Such a finding implies that inserting a needle into the left hand would potentially lead to pain relief in the right hand. The recent findings of Hsieh *et al.*³³ add support to the practice of needling contralateral to the side of pain and/or needling distal to the site of pain in order to reduce end plate noise and suppress activity of more proximal painful TrPs.

In another recent experimental study,²⁶ dry needling of primary MTrPs in patients with shoulder pain was found to inhibit the activity of satellite or secondary MTrPs situated in the shoulder, elbow, or forearm. That is, the target MTrP was found either proximal or distal to the patient’s site of discomfort.²⁶ It follows then that the APTA⁷ and Physical Therapy State Boards^{10,13} should consider that the best practice for dry needling may require the placement of needles both ‘local and/or distal’ to the patient’s primary source of pain.

Physical therapists often deliver manual therapy and/or orthotic interventions to the tarso-metatarsal, subtalar, or talocrural joints in order to manage patients with low back pain;¹⁹²⁻¹⁹⁶ likewise, the insertion of needles without injectate into bodily areas that are asymptomatic but distal or proximal to the site of pain is supported by the myofascial pain syndrome literature.¹⁵⁴ As Melzack *et al.*²⁷ points out:

The stimulation of particular nerves or tissues by needles...would close the gates to [pain] inputs from selected body areas. The cells of the midbrain reticular formation are known to have large receptive fields.... It is possible, then, that particular [distant] body areas may project especially strongly to some reticular areas, and these, in turn, could bring about a complete block of inputs from particular parts of the body.²⁷

Conclusion

Just as the Maitland,¹⁴ Kaltenborn,¹⁵ and McKenzie¹⁶ approaches are unique brands of manual therapy, so is 'TDN' and 'IMT' to dry needling. They provide a single framework or paradigm within the much broader field of dry needling.^{5,17–25} Therefore, the terms 'TDN' or 'IMT' should not be used interchangeably or synonymously with the term 'Dry Needling'. Dry needling encompasses the insertion of needles without injectate³ into, alongside, or around nerves,^{17,28,36,40,42,43,47,55,65,164} muscles,^{5,6,20,23,24,65,109} or connective tissues^{6,7,48,49,70–72,126,170,172} for the management of pain and dysfunction in neuromusculoskeletal conditions. Dry needling neither attempts to move *qi* along meridians nor does it rely on diagnoses from traditional Chinese or Oriental Medicine.^{78,79} Considering the mechanical,^{70–72,163} chemical,^{146,164} endocrinological,¹³⁶ microvascular,^{138,164,165,167,168,171} neural,^{36,41,47} and central effects^{38,39,44,46,130,137,140} of both superficial and deep dry needling at TrP and non-TrP locations, the position statements by the APTA^{8,75} and many State Boards of Physical Therapy^{9–13} do not seem to be consistent with the existing literature.

References

- Speed CA. Injection therapies for soft-tissue disorders. *Best Pract Res Clin Rheumatol.* 2003;**17**(1):167–81.
- Speed CA. Injection therapies for soft-tissue lesions. *Best Pract Res Clin Rheumatol.* 2007;**21**(2):333–47.
- Casanueva B, Rivas P, Rodero B, Quintal C, Llorca J, Gonzalez-Gay MA. Short-term improvement following dry needle stimulation of tender points in fibromyalgia. *Rheumatol Int.* Apr 23, 2013. [epub ahead of print]
- Manheimer E, White A, Berman B, Forsys K, Ernst E. Meta-analysis: acupuncture for low back pain. *Ann Intern Med.* 2005;**142**(8):651–63.
- Gunn CC, Milbrandt WE, Little AS, Mason KE. Dry needling of muscle motor points for chronic low-back pain: a randomized clinical trial with long-term follow-up. *Spine (Phila Pa 1976).* 1980;**5**(3):279–91.
- Lewit K. The needle effect in the relief of myofascial pain. *Pain.* 1979;**6**(1):83–90.
- Neal BS, Longbottom J. Is there a role for acupuncture in the treatment of tendinopathy? *Acupunct Med.* 2012;**30**(4):346–9.
- APTA. Physical therapists & the performance of dry needling: an educational resource paper. Alexandria, VA, USA: APTA Department of Practice and APTA State Government Affairs; 2012.
- NCBPTE. Position statement: intramuscular manual therapy (dry needling). North Carolina Board of Physical Therapy Examiners, June 14, 2012.
- MSSBPT. Intramuscular manual therapy (dry needling) may be performed by a licensed physical therapist. Mississippi State Board of Physical Therapy, Part 3101 Rule 1.3c, September 10, 2012.
- LAPTB. Louisiana Physical Therapy Practice Act: Treatment with Dry Needling. Louisiana Physical Therapy Board, Rule 311, Page 29, October 20, 2011.
- NEBPT. Minutes of the board of physical therapy: discussion of board opinions. Nebraska Board of Physical Therapy, June 20, 2011:1–4.
- SPTB. Requirements for physical therapists to perform dry needling. Colorado Department of Regulatory Agencies: State Physical Therapy Board, Rule 211, 2008:10–11.
- Maitland GD. *Vertebral manipulation*, 2nd edn. London: Butterworths; 1968.
- Kaltenborn FM. *Mobilization of the spine*, 1st edn. Oslo, Norway: Olaf Norlis Bokhandel; 1970.
- McKenzie R, May S. *The cervical and thoracic spine: mechanical diagnosis and therapy*, 2nd edn, Vol. 1. Waikanae, NZ: Spinal Publications New Zealand; 2006.
- Cummings TM, White AR. Needling therapies in the management of myofascial trigger point pain: a systematic review. *Arch Phys Med Rehabil.* 2001;**82**(7):986–92.
- DiLorenzo L, Trallesi M, Morelli D, Pompa A, Brunelli S, Buzzi MG, et al. Hemiparetic shoulder pain syndrome treated with deep dry needling during early rehabilitation: a prospective, open-label, randomized investigation. *J Musculoskeletal Pain.* 2004;**12**(2):25–34.
- Furlan AD, van Tulder M, Cherkin D, Tsukayama H, Lao L, Koes B, et al. Acupuncture and dry-needling for low back pain: an updated systematic review within the framework of the cochrane collaboration. *Spine (Phila Pa 1976).* 2005;**30**(8):944–63.
- Itoh K, Katsumi Y, Kitakoji H. Trigger point acupuncture treatment of chronic low back pain in elderly patients—a blinded RCT. *Acupunct Med.* 2004;**22**(4):170–7.
- Kalichman L, Vulfsons S. Dry needling in the management of musculoskeletal pain. *J Am Board Fam Med.* 2010;**23**(5):640–6.
- Myburgh C, Larsen AH, Hartvigsen J. A systematic, critical review of manual palpation for identifying myofascial trigger points: evidence and clinical significance. *Arch Phys Med Rehabil.* 2008;**89**(6):1169–76.
- Tough EA, White A. Effectiveness of acupuncture/dry needling for myofascial trigger point pain—a systematic review. *Phys Ther Rev.* 2011;**16**(2):147–54.
- Tough EA, White AR, Cummings TM, Richards SH, Campbell JL. Acupuncture and dry needling in the management of myofascial trigger point pain: a systematic review and meta-analysis of randomised controlled trials. *Eur J Pain.* 2009;**13**(1):3–10.
- Tough EA, White AR, Richards S, Campbell J. Variability of criteria used to diagnose myofascial trigger point pain syndrome—evidence from a review of the literature. *Clin J Pain.* 2007;**23**(3):278–86.
- Hsieh YL, Kao MJ, Kuan TS, Chen SM, Chen JT, Hong CZ. Dry needling to a key myofascial trigger point may reduce the irritability of satellite MTrPs. *Am J Phys Med Rehabil.* 2007;**86**(5):397–403.
- Melzack R, Stillwell DM, Fox EJ. Trigger points and acupuncture points for pain: correlations and implications. *Pain.* 1977;**3**(1):3–23.
- Hong C. Treatment of myofascial pain syndrome. *Curr Pain Headache Rep.* 2006;**10**:345–9.
- Hong CZ. Lidocaine injection versus dry needling to myofascial trigger point. The importance of the local twitch response. *Am J Phys Med Rehabil.* 1994;**73**(4):256–63.
- Hong CZ, Kuan TS, Chen JT, Chen SM. Referred pain elicited by palpation and by needling of myofascial trigger points: a comparison. *Arch Phys Med Rehabil.* 1997;**78**(9):957–60.
- Hong C. Myofascial trigger points: pathophysiology and correlation with acupuncture points. *Acupunct Med.* 2000;**18**(1):41–7.
- Hong CZ, Simons DG. Pathophysiologic and electrophysiologic mechanisms of myofascial trigger points. *Arch Phys Med Rehabil.* 1998;**79**(7):863–72.
- Hsieh YL, Chou LW, Joe YS, Hong CZ. Spinal cord mechanism involving the remote effects of dry needling on the irritability of myofascial trigger spots in rabbit skeletal muscle. *Arch Phys Med Rehabil.* 2011;**92**(7):1098–105.
- Audette JF, Wang F, Smith H. Bilateral activation of motor unit potentials with unilateral needle stimulation of active myofascial trigger points. *Am J Phys Med Rehabil.* 2004;**83**(5):368–74.
- Kamanli A, Kaya A, Ardicoglu O, Ozgocmen S, Zengin FO, Bayik Y. Comparison of lidocaine injection, botulinum toxin injection, and dry needling to trigger points in myofascial pain syndrome. *Rheumatol Int.* 2005;**25**(8):604–11.
- Khosrawi S, Moghtaderi A, Haghghat S. Acupuncture in treatment of carpal tunnel syndrome: a randomized controlled trial study. *J Res Med Sci.* 2012;**17**(1):1–7.
- Kummerdee W, Kaewtong A. Efficacy of acupuncture versus night splinting for carpal tunnel syndrome: a randomized clinical trial. *J Med Assoc Thai.* 2010;**93**(12):1463–9.
- Hsieh JC, Tu CH, Chen FP, Chen MC, Yeh TC, Cheng HC, et al. Activation of the hypothalamus characterizes the acupuncture stimulation at the analgesic point in human: a

- positron emission tomography study. *Neurosci Lett*. 2001;**307**(2):105–8.
- 39 Hui KK, Liu J, Makris N, Gollub RL, Chen AJ, Moore CI, et al. Acupuncture modulates the limbic system and sub-cortical gray structures of the human brain: evidence from fMRI studies in normal subjects. *Hum Brain Mapp*. 2000;**9**(1):13–25.
 - 40 Yang CP, Hsieh CL, Wang NH, Li TC, Hwang KL, Yu SC, et al. Acupuncture in patients with carpal tunnel syndrome: a randomized controlled trial. *Clin J Pain*. 2009;**25**(4):327–33.
 - 41 Yang CP, Wang NH, Li TC, Hsieh CL, Chang HH, Hwang KL, et al. A randomized clinical trial of acupuncture versus oral steroids for carpal tunnel syndrome: a long-term follow-up. *J Pain*. 2011;**12**(2):272–9.
 - 42 Ma YT. Biomedical acupuncture for sports and trauma rehabilitation: dry needling techniques. St. Louis, MO, USA: Churchill Livingstone/Elsevier; 2011.
 - 43 Ma YT, Ma M, Cho ZH. Biomedical acupuncture for pain management: an integrative approach. St. Louis, MO, USA: Elsevier Churchill Livingstone; 2005.
 - 44 Napadow V, Kettner N, Liu J, Li M, Kwong KK, Vangel M, et al. Hypothalamus and amygdala response to acupuncture stimuli in Carpal Tunnel Syndrome. *Pain*. 2007;**130**(3):254–66.
 - 45 Napadow V, Kettner N, Ryan A, Kwong KK, Audette J, Hui KK. Somatosensory cortical plasticity in carpal tunnel syndrome—a cross-sectional fMRI evaluation. *Neuroimage*. 2006;**31**(2):520–30.
 - 46 Napadow V, Makris N, Liu J, Kettner NW, Kwong KK, Hui KK. Effects of electroacupuncture versus manual acupuncture on the human brain as measured by fMRI. *Hum Brain Mapp*. 2005;**24**(3):193–205.
 - 47 Sim H, Shin BC, Lee MS, Jung A, Lee H, Ernst E. Acupuncture for carpal tunnel syndrome: a systematic review of randomized controlled trials. *J Pain*. 2011;**12**(3):307–14.
 - 48 Imamura M, Fischer A, Imamura S, Kaziyama H, Carvalho A, Salomao O. Treatment of myofascial pain components in plantar fasciitis speeds up recovery: documentation by algometry. *J Musculoskel Pain*. 1998;**6**(1):91–110.
 - 49 Tillu A, Gupta S. Effect of acupuncture treatment on heel pain due to plantar fasciitis. *Acupunct Med*. 1998;**16**(2):66–8.
 - 50 Vas J, White A. Evidence from RCTs on optimal acupuncture treatment for knee osteoarthritis—an exploratory review. *Acupunct Med*. 2007;**25**(1–2):29–35.
 - 51 White A, Foster NE, Cummings M, Barlas P. Acupuncture treatment for chronic knee pain: a systematic review. *Rheumatology (Oxford)*. 2007;**46**(3):384–90.
 - 52 Whitehurst DG, Bryan S, Hay EM, Thomas E, Young J, Foster NE. Cost-effectiveness of acupuncture care as an adjunct to exercise-based physical therapy for osteoarthritis of the knee. *Phys Ther*. 2011;**91**(5):630–41.
 - 53 Witt CM, Jena S, Brinkhaus B, Liecker B, Wegscheider K, Willich SN. Acupuncture in patients with osteoarthritis of the knee or hip: a randomized, controlled trial with an additional nonrandomized arm. *Arthritis Rheum*. 2006;**54**(11):3485–93.
 - 54 Zhang S, Yip T, Li Q. Acupuncture treatment for plantar fasciitis: a randomized controlled trial with six months follow-up. *Evid Based Complement Alternat Med*. 2009;**20**(11):1–10.
 - 55 Baldry P. Acupuncture, trigger points, and musculoskeletal pain. Edinburgh, New York, NY, USA: Churchill Livingstone; 1989.
 - 56 Cotchett MP, Landorf KB, Munteanu SE. Effectiveness of dry needling and injections of myofascial trigger points associated with plantar heel pain: a systematic review. *J Foot Ankle Res*. 2010;**3**:18.
 - 57 Dorsher PT, Fleckenstein J. Trigger points and classical acupuncture points: part 1: qualitative and quantitative anatomic correspondences. *Ger J Acupunct Relat Tech*. 2008;**51**(3):15–24.
 - 58 Dorsher PT, Fleckenstein J. Trigger points and classical acupuncture points: part 2: qualitative correspondences in treating pain and somatovisceral disorders. *Ger J Acupunct Relat Tech*. 2008;**51**(4):6–11.
 - 59 Fernandez-Carnero J, La Touche R, Ortega-Santiago R, Galan-del-Rio F, Pesquera J, Ge HY, et al. Short-term effects of dry needling of active myofascial trigger points in the masseter muscle in patients with temporomandibular disorders. *J Orofac Pain*. 2010;**24**(1):106–12.
 - 60 Fishman LM, Dombi GW, Michaelsen C, Ringel S, Rozbruch J, Rosner B, et al. Piriformis syndrome: diagnosis, treatment, and outcome—a 10-year study. *Arch Phys Med Rehabil*. 2002;**83**(3):295–301.
 - 61 Franca DL, Senna-Fernandes V, Cortez CM, Jackson MN, Bernardo-Filho M, Guimaraes MA. Tension neck syndrome treated by acupuncture combined with physiotherapy: a comparative clinical trial (pilot study). *Complement Ther Med*. 2008;**16**(5):268–77.
 - 62 Furlan AD, van Tulder MW, Cherkin DC, Tsukayama H, Lao L, Koes BW, et al. Acupuncture and dry-needling for low back pain. *Cochrane Database Syst Rev*. 2005;(1):CD001351.
 - 63 Ga H, Choi JH, Park CH, Yoon HJ. Acupuncture needling versus lidocaine injection of trigger points in myofascial pain syndrome in elderly patients—a randomised trial. *Acupunct Med*. 2007;**25**(4):130–6.
 - 64 Ga H, Choi JH, Park CH, Yoon HJ. Dry needling of trigger points with and without paraspinal needling in myofascial pain syndromes in elderly patients. *J Altern Complement Med*. 2007;**13**(6):617–24.
 - 65 Ga H, Koh HJ, Choi JH, Kim CH. Intramuscular and nerve root stimulation vs lidocaine injection to trigger points in myofascial pain syndrome. *J Rehabil Med*. 2007;**39**(5):374–8.
 - 66 Garvey TA, Marks MR, Wiesel SW. A prospective, randomized, double-blind evaluation of trigger-point injection therapy for low-back pain. *Spine (Phila Pa 1976)*. 1989;**14**(9):962–4.
 - 67 Gonzalez-Perez LM, Infante-Cossio P, Granados-Nunez M, Urresti-Lopez FJ. Treatment of temporomandibular myofascial pain with deep dry needling. *Med Oral Patol Oral Cir Bucal*. 2012;**17**(5):e781–5.
 - 68 Gunn CC. The Gunn approach to the treatment of chronic pain: intramuscular stimulation for myofascial pain of radiculopathic origin, 2nd edn. New York, NY, USA: Churchill Livingstone; 1996.
 - 69 Haake M, Muller HH, Schade-Brittinger C, Schäfer H, Maier C, Endres HG, et al. German Acupuncture Trials (GERAC) for chronic low back pain: randomized, multicenter, blinded, parallel-group trial with 3 groups. *Arch Intern Med*. 2007;**167**(17):1892–8.
 - 70 Langevin HM, Bouffard NA, Badger GJ, Churchill DL, Howe AK. Subcutaneous tissue fibroblast cytoskeletal remodeling induced by acupuncture: evidence for a mechanotransduction-based mechanism. *J Cell Physiol*. 2006;**207**(3):767–74.
 - 71 Langevin HM, Bouffard NA, Badger GJ, Iatridis JC, Howe AK. Dynamic fibroblast cytoskeletal response to subcutaneous tissue stretch *ex vivo* and *in vivo*. *Am J Physiol Cell Physiol*. 2005;**288**(3):C747–56.
 - 72 Langevin HM, Churchill DL, Cipolla MJ. Mechanical signaling through connective tissue: a mechanism for the therapeutic effect of acupuncture. *Faseb J*. 2001;**15**(12):2275–82.
 - 73 Travell JG, Simons DG. Myofascial pain and dysfunction: the trigger point manual, Vol. 1. Baltimore, MD, USA: Williams and Wilkins; 1983.
 - 74 Travell JG, Simons DG. Myofascial pain and dysfunction: the trigger point manual, Vol. 2. Baltimore, MD, USA: Williams and Wilkins; 1992.
 - 75 APTA. Description of dry needling in clinical practice: an educational resource paper. Alexandria, VA, USA: APTA Public Policy, Practice, and Professional Affairs Unit; 2013.
 - 76 AZPTA. Definition and position statement for dry needling by physical therapists in Arizona. Arizona Physical Therapy Association, 2013.
 - 77 Rathnayake T. Back pain (low): acupuncture and dry needling. Evidence Summaries-Joanna Briggs Institute; 2009.
 - 78 Deadman P, Al-Khafaji M, Baker K. A manual of acupuncture, 2nd ed. Hove, East Sussex, UK: Journal of Chinese Medicine Publications; 2011.
 - 79 O'Conner J, Bensky D. Acupuncture: a comprehensive text. Seattle, WA, USA: Eastland Press; 1981.
 - 80 Itoh K, Katsumi Y, Hirota S, Kitakoji H. Randomised trial of trigger point acupuncture compared with other acupuncture for treatment of chronic neck pain. *Complement Ther Med*. 2007;**15**(3):172–9.
 - 81 Berman BM, Lao L, Langenberg P, Lee WL, Gilpin AM, Hochberg MC. Effectiveness of acupuncture as adjunctive therapy in osteoarthritis of the knee: a randomized, controlled trial. *Ann Intern Med*. 2004;**141**(12):901–10.
 - 82 Berman BM, Singh BB, Lao L, Langenberg P, Li H, Hadhazy V, et al. A randomized trial of acupuncture as an adjunctive therapy in osteoarthritis of the knee. *Rheumatology (Oxford)*. 1999;**38**(4):346–54.

- 83 Ezzo J, Hadhazy V, Birch S, Lao L, Kaplan G, Hochberg M, *et al.* Acupuncture for osteoarthritis of the knee: a systematic review. *Arthritis Rheum.* 2001;**44**(4):819–25.
- 84 Foster NE, Thomas E, Barlas P, Hill JC, Young J, Mason E, *et al.* Acupuncture as an adjunct to exercise based physiotherapy for osteoarthritis of the knee: randomised controlled trial. *BMJ.* 2007;**335**(7617):436.
- 85 Kwon YD, Pittler MH, Ernst E. Acupuncture for peripheral joint osteoarthritis: a systematic review and meta-analysis. *Rheumatology (Oxford).* 2006;**45**(11):1331–7.
- 86 Manheimer E, Linde K, Lao L, Bouter LM, Berman BM. Meta-analysis: acupuncture for osteoarthritis of the knee. *Ann Intern Med.* 2007;**146**(12):868–77.
- 87 Mavrommatis CI, Argyra E, Vadalouka A, Vasilakos DG. Acupuncture as an adjunctive therapy to pharmacological treatment in patients with chronic pain due to osteoarthritis of the knee: a 3-armed, randomized, placebo-controlled trial. *Pain.* 2012;**153**(8):1720–6.
- 88 Scharf HP, Mansmann U, Streitberger K, Witte S, Krämer J, Maier C, *et al.* Acupuncture and knee osteoarthritis: a three-armed randomized trial. *Ann Intern Med.* 2006;**145**(1):12–20.
- 89 Vas J, Mendez C, Perea-Milla E, Vega E, Panadero MD, León JM, *et al.* Acupuncture as a complementary therapy to the pharmacological treatment of osteoarthritis of the knee: randomised controlled trial. *BMJ.* 2004;**329**(7476):1216.
- 90 Witt C, Brinkhaus B, Jena S, Linde K, Streng A, Wagenpfeil S, *et al.* Acupuncture in patients with osteoarthritis of the knee: a randomised trial. *Lancet.* 2005;**366**(9480):136–43.
- 91 Fink MG, Kunsebeck H, Wipperman B, Gehrke A. Non-specific effects of traditional Chinese acupuncture in osteoarthritis of the hip. *Complement Ther Med.* 2001;**9**(2):82–9.
- 92 Haslam R. A comparison of acupuncture with advice and exercises on the symptomatic treatment of osteoarthritis of the hip—a randomised controlled trial. *Acupunct Med.* 2001;**19**(1):19–26.
- 93 Stener-Victorin E, Kruse-Smidje C, Jung K. Comparison between electro-acupuncture and hydrotherapy, both in combination with patient education and patient education alone, on the symptomatic treatment of osteoarthritis of the hip. *Clin J Pain.* 2004;**20**(3):179–85.
- 94 Zhang W, Moskowitz RW, Nuki G, Abramson S, Altman RD, Arden N, *et al.* OARSI recommendations for the management of hip and knee osteoarthritis, Part II: OARSI evidence-based, expert consensus guidelines. *Osteoarthritis Cartilage.* 2008;**16**(2):137–62.
- 95 Allais G, De Lorenzo C, Quirico PE, Airola G, Tolardo G, Mana O, *et al.* Acupuncture in the prophylactic treatment of migraine without aura: a comparison with flunarizine. *Headache.* 2002;**42**(9):855–61.
- 96 Coeytaux RR, Kaufman JS, Kaptchuk TJ, Chen W, Miller WC, Callahan LF, *et al.* A randomized, controlled trial of acupuncture for chronic daily headache. *Headache.* 2005;**45**(9):1113–23.
- 97 Melchart D, Linde K, Fischer P, White A, Allais G, Vickers A, *et al.* Acupuncture for recurrent headaches: a systematic review of randomized controlled trials. *Cephalalgia.* 1999;**19**(9):779–86; discussion 765.
- 98 Melchart D, Linde K, Streng A, Reitmayr S, Hoppe A, Brinkhaus B, Becker-Witt C, *et al.* Acupuncture Randomized Trials (ART) in patients with migraine or tension-type headache—design and protocols. *Forsch Komplementarmed Klass Naturheilkd.* 2003;**10**(4):179–84.
- 99 Melchart D, Thormaehlen J, Hager S, Liao J, Linde K, Weidenhammer W. Acupuncture versus placebo versus sumatriptan for early treatment of migraine attacks: a randomized controlled trial. *J Intern Med.* 2003;**253**(2):181–8.
- 100 Vickers AJ, Rees RW, Zollman CE, McCarney R, Smith CM, Ellis N, *et al.* Acupuncture for chronic headache in primary care: large, pragmatic, randomised trial. *BMJ.* 2004;**328**(7442):744.
- 101 Wonderling D, Vickers AJ, Grieve R, McCarney R. Cost effectiveness analysis of a randomised trial of acupuncture for chronic headache in primary care. *BMJ.* 2004;**328**(7442):747.
- 102 Venancio Rde A, Alencar FG Jr, Zamperini C. Botulinum toxin, lidocaine, and dry-needling injections in patients with myofascial pain and headaches. *Cranio.* 2009;**27**(1):46–53.
- 103 Manolopoulos L, Vlastarakos PV, Georgiou L, Giotakis I, Loizos A, Nikolopoulos TP. Myofascial pain syndromes in the maxillofacial area: a common but underdiagnosed cause of head and neck pain. *Int J Oral Maxillofac Surg.* 2008;**37**(11):975–84.
- 104 McMillan AS, Nolan A, Kelly PJ. The efficacy of dry needling and procaine in the treatment of myofascial pain in the jaw muscles. *J Orofac Pain.* 1997;**11**(4):307–14.
- 105 Shen YF, Goddard G. The short-term effects of acupuncture on myofascial pain patients after clenching. *Pain Pract.* 2007;**7**(3):256–64.
- 106 Shen YF, Younger J, Goddard G, Mackey S. Randomized clinical trial of acupuncture for myofascial pain of the jaw muscles. *J Orofac Pain.* 2009;**23**(4):353–9.
- 107 Smith P, Mossrop D, Davies S, Sloan P, Al-Ani Z. The efficacy of acupuncture in the treatment of temporomandibular joint myofascial pain: a randomised controlled trial. *J Dent.* 2007;**35**(3):259–67.
- 108 Green S, Buchbinder R, Hetrick S. Acupuncture for shoulder pain. *Cochrane Database Syst Rev.* 2005;(2):CD005319.
- 109 Ingber RS. Shoulder impingement in tennis/racquetball players treated with subscapularis myofascial treatments. *Arch Phys Med Rehabil.* 2000;**81**(5):679–82.
- 110 Osborne NJ, Gatt IT. Management of shoulder injuries using dry needling in elite volleyball players. *Acupunct Med.* 2010;**28**(1):42–5.
- 111 Irnich D, Behrens N, Gleditsch JM, Stör W, Schreiber MA, Schöps P, *et al.* Immediate effects of dry needling and acupuncture at distant points in chronic neck pain: results of a randomized, double-blind, sham-controlled crossover trial. *Pain.* 2002;**99**(1–2):83–9.
- 112 Irnich D, Behrens N, Molzen H, König A, Gleditsch J, Krauss M, *et al.* Randomised trial of acupuncture compared with conventional massage and ‘sham’ laser acupuncture for treatment of chronic neck pain. *BMJ.* 2001;**322**(7302):1574–8.
- 113 Ma C, Wu S, Li G, Xiao X, Mai M, Yan T. Comparison of miniscalpel-needle release, acupuncture needling, and stretching exercise to trigger point in myofascial pain syndrome. *Clin J Pain.* 2010;**26**(3):251–7.
- 114 Trinh K, Graham N, Gross A, Goldsmith C, Wang E, Cameron I, *et al.* Acupuncture for neck disorders. *Spine (Phila Pa 1976).* 2007;**32**(2):236–43.
- 115 Brinkhaus B, Witt CM, Jena S, Linde K, Streng A, Wagenpfeil S, *et al.* Acupuncture in patients with chronic low back pain: a randomized controlled trial. *Arch Intern Med.* 2006;**166**(4):450–7.
- 116 Cherkin DC, Sherman KJ, Avins AL, Erro JH, Ichikawa L, Barlow WE, *et al.* A randomized trial comparing acupuncture, simulated acupuncture, and usual care for chronic low back pain. *Arch Intern Med.* 2009;**169**(9):858–66.
- 117 Chou R, Qaseem A, Snow V, Casey D, Cross JT Jr, Shekelle P, *et al.* Diagnosis and treatment of low back pain: a joint clinical practice guideline from the American College of Physicians and the American Pain Society. *Ann Intern Med.* 2007;**147**(7):478–91.
- 118 Eisenberg DM, Post DE, Davis RB, Connelly MT, Legedza AT, Hrbek AL, *et al.* Addition of choice of complementary therapies to usual care for acute low back pain: a randomized controlled trial. *Spine (Phila Pa 1976).* 2007;**32**(2):151–8.
- 119 Giles LG, Muller R. Chronic spinal pain: a randomized clinical trial comparing medication, acupuncture, and spinal manipulation. *Spine (Phila Pa 1976).* 2003;**28**(14):1490–502; discussion 1502–493.
- 120 Inoue M, Kitakoji H, Ishizaki N, Tawa M, Yano T, Katsumi Y, *et al.* Relief of low back pain immediately after acupuncture treatment—a randomised, placebo controlled trial. *Acupunct Med.* 2006;**24**(3):103–8.
- 121 Johnston BC, da Costa BR, Devereaux PJ, Akl EA, Busse JW. The use of expertise-based randomized controlled trials to assess spinal manipulation and acupuncture for low back pain: a systematic review. *Spine (Phila Pa 1976).* 2008;**33**(8):914–8.
- 122 MacPherson H, Thorpe L, Thomas K, Campbell M. Acupuncture for low back pain: traditional diagnosis and treatment of 148 patients in a clinical trial. *Complement Ther Med.* 2004;**12**(1):38–44.
- 123 Molsberger AF, Mau J, Pawelec DB, Winkler J. Does acupuncture improve the orthopedic management of chronic low back pain—a randomized, blinded, controlled trial with 3 months follow up. *Pain.* 2002;**99**(3):579–87.
- 124 Savigny P, Watson P, Underwood M. Early management of persistent non-specific low back pain: summary of NICE guideline. *Br Med J.* 2009;**338**:1441–5.
- 125 Vas J, Aranda JM, Modesto M, Benítez-Parejo N, Herrera A, Martínez-Barquín DM, *et al.* Acupuncture in patients with acute low back pain: a multicentre randomised controlled clinical trial. *Pain.* 2012;**153**(9):1883–9.

- 126 Ebrahim AHM, Ahmed GM, Elsayed E, Sarhan R. Effect of electroacupuncture TENS, stretching exercises, and prefabricated insole in patients with plantar fasciitis. *Sci J Al-Azhar Med Univ.* 2007;**28**(3):1–10.
- 127 Perez-Millan R, Foster L. Low-frequency electroacupuncture in the management of refractory plantar fasciitis: a case-series. *Med Acupunct.* 2001;**13**(1):1–7.
- 128 Baldry P. The integration of acupuncture within medicine in the UK—the British Medical Acupuncture Society's 25th anniversary. *Acupunct Med.* 2005;**23**(1):2–12.
- 129 Lee SH, Chen CC, Lee CS, Lin TC, Chan RC. Effects of needle electrical intramuscular stimulation on shoulder and cervical myofascial pain syndrome and microcirculation. *J Chin Med Assoc.* 2008;**71**(4):200–6.
- 130 Biella G, Sotgiu ML, Pellegata G, Paulesu E, Castiglioni I, Fazio F. Acupuncture produces central activations in pain regions. *Neuroimage.* 2001;**14**(1 Pt 1):60–6.
- 131 Foster NE, Thomas E, Hill JC, Hay EM. The relationship between patient and practitioner expectations and preferences and clinical outcomes in a trial of exercise and acupuncture for knee osteoarthritis. *Eur J Pain.* 2010;**14**(4):402–9.
- 132 White P, Lewith G, Prescott P, Conway J. Acupuncture versus placebo for the treatment of chronic mechanical neck pain: a randomized, controlled trial. *Ann Intern Med.* 2004;**141**(12):911–9.
- 133 Witt CM, Jena S, Brinkhaus B, Liecker B, Wegscheider K, Willich SN. Acupuncture for patients with chronic neck pain. *Pain.* 2006;**125**(1–2):98–106.
- 134 Witt CM, Jena S, Selim D, Brinkhaus B, Reinhold T, Wruck K, et al. Pragmatic randomized trial evaluating the clinical and economic effectiveness of acupuncture for chronic low back pain. *Am J Epidemiol.* 2006;**164**(5):487–96.
- 135 Yuan J, Purepong N, Kerr DP, Park J, Bradbury I, McDonough S. Effectiveness of acupuncture for low back pain: a systematic review. *Spine (Phila Pa 1976).* 2008;**33**(23):E887–900.
- 136 Ahsin S, Saleem S, Bhatti AM, Iles RK, Aslam M. Clinical and endocrinological changes after electro-acupuncture treatment in patients with osteoarthritis of the knee. *Pain.* 2009;**147**(1–3):60–6.
- 137 Almeida RT, Perez AC, Francischi JN, Castro MS, Duarte ID. Opioidergic orofacial antinociception induced by electroacupuncture at acupoint St36. *Braz J Med Biol Res.* 2008;**41**(7):621–6.
- 138 Loaiza LA, Yamaguchi S, Ito M, Ohshima N. Electroacupuncture stimulation to muscle afferents in anesthetized rats modulates the blood flow to the knee joint through autonomic reflexes and nitric oxide. *Auton Neurosci.* 2002;**97**(2):103–9.
- 139 Yeung CK, Leung MC, Chow DH. The use of electro-acupuncture in conjunction with exercise for the treatment of chronic low-back pain. *J Altern Complement Med.* 2003;**9**(4):479–90.
- 140 Almeida RT, Duarte ID. Nitric oxide/cGMP pathway mediates orofacial antinociception induced by electroacupuncture at the St36 acupoint. *Brain Res.* 2008;**1188**:54–60.
- 141 Ceccherelli F, Gioioso L, Casale R, Gagliardi G, Ori C. Neck pain treatment with acupuncture: does the number of needles matter? *Clin J Pain.* 2010;**26**(9):807–12.
- 142 Bovey M. Deqi. *J Chin Med.* 2006;**81**:18–29.
- 143 Hui KK, Nixon EE, Vangel MG, Liu J, Marina O, Napadow V, et al. Characterization of the 'deqi' response in acupuncture. *BMC Complement Altern Med.* 2007;**7**:33.
- 144 Thomas KJ, MacPherson H, Thorpe L, Brazier J, Fitter M, Campbell MJ, et al. Randomised controlled trial of a short course of traditional acupuncture compared with usual care for persistent non-specific low back pain. *BMJ.* 2006;**333**(7569):623.
- 145 Rha DW, Shin JC, Kim YK, Jung JH, Kim YU, Lee SC. Detecting local twitch responses of myofascial trigger points in the lower-back muscles using ultrasonography. *Arch Phys Med Rehabil.* 2011;**92**(10):1576–80.e1.
- 146 Shah JP, Danoff JV, Desai MJ, Parikh S, Nakamura LY, Phillips TM, et al. Biochemicals associated with pain and inflammation are elevated in sites near to and remote from active myofascial trigger points. *Arch Phys Med Rehabil.* 2008;**89**(1):16–23.
- 147 Sciotti VM, Mittak VL, DiMarco L, Ford LM, Plezbert J, Santipadri E, et al. Clinical precision of myofascial trigger point location in the trapezius muscle. *Pain.* 2001;**93**(3):259–66.
- 148 Brinkhaus B, Witt C, Jena S, Linde K, Streng A, Irnich D, et al. Interventions and physician characteristics in a randomized multicenter trial of acupuncture in patients with low-back pain. *J Altern Complement Med.* 2006;**12**(7):649–57.
- 149 Lucas N, Macaskill P, Irwig L, Moran R, Bogduk N. Reliability of physical examination for diagnosis of myofascial trigger points: a systematic review of the literature. *Clin J Pain.* 2009;**25**(1):80–9.
- 150 Trinh KV, Graham N, Gross AR, Goldsmith CH, Wang E, Cameron ID, et al. Acupuncture for neck disorders. *Cochrane Database Syst Rev.* 2006;(3):CD004870.
- 151 Airaksinen O, Brox JI, Cedraschi C, Hildebrandt J, Klaber-Moffett J, Kovacs F, et al. Chapter 4. European guidelines for the management of chronic nonspecific low back pain. *Eur Spine J.* 2006;**15**(Suppl 2):S192–300.
- 152 van Tulder M, Becker A, Bekkering T, Breen A, del Real MT, Hutchinson A, et al. Chapter 3. European guidelines for the management of acute nonspecific low back pain in primary care. *Eur Spine J.* 2006;**15**(Suppl 2):S169–191.
- 153 Moffet H. Traditional acupuncture theories yield null outcomes: a systematic review of clinical trials. *J Clin Epidemiol.* 2008;**61**:741–7.
- 154 Dorsher PT. Myofascial referred-pain data provide physiologic evidence of acupuncture meridians. *J Pain.* 2009;**10**(7):723–31.
- 155 Barbero M, Bertoli P, Cescon C, Macmillan F, Couto F, Gatti R. Intra-rater reliability of an experienced physiotherapist in locating myofascial trigger points in upper trapezius muscle. *J Man Manip Ther.* 2012;**20**:171–7.
- 156 Lew PC, Lewis J, Story I. Inter-therapist reliability in locating latent myofascial trigger points using palpation. *Man Ther.* 1997;**2**(2):87–90.
- 157 Myburgh C, Lauridsen HH, Larsen AH, Hartvigsen J. Standardized manual palpation of myofascial trigger points in relation to neck/shoulder pain; the influence of clinical experience on inter-examiner reproducibility. *Man Ther.* 2011;**16**(2):136–40.
- 158 Al-Shenqiti AM, Oldham JA. Test-retest reliability of myofascial trigger point detection in patients with rotator cuff tendonitis. *Clin Rehabil.* 2005;**19**(5):482–7.
- 159 Gerwin RD, Shannon S, Hong CZ, Hubbard D, Gevirtz R. Interrater reliability in myofascial trigger point examination. *Pain.* 1997;**69**(1–2):65–73.
- 160 Hsieh CY, Hong CZ, Adams AH, Platt KJ, Danielson CD, Hoehler FK, et al. Interexaminer reliability of the palpation of trigger points in the trunk and lower limb muscles. *Arch Phys Med Rehabil.* 2000;**81**(3):258–64.
- 161 Levoska S, Keinanen-Kiukaanniemi S, Bloigu R. Repeatability of measurement of tenderness in the neck-shoulder region by a dolorimeter and manual palpation. *Clin J Pain.* 1993;**9**(4):229–35.
- 162 Njoo KH, Van der Does E. The occurrence and inter-rater reliability of myofascial trigger points in the quadratus lumborum and gluteus medius: a prospective study in non-specific low back pain patients and controls in general practice. *Pain.* 1994;**58**(3):317–23.
- 163 Langevin HM, Bouffard NA, Churchill DL, Badger GJ. Connective tissue fibroblast response to acupuncture: dose-dependent effect of bidirectional needle rotation. *J Altern Complement Med.* 2007;**13**(3):355–60.
- 164 Shah JP, Phillips TM, Danoff JV, Gerber LH. An *in vivo* microanalytical technique for measuring the local biochemical milieu of human skeletal muscle. *J Appl Physiol.* 2005;**99**(5):1977–84.
- 165 Kubo K, Yajima H, Takayama M, Ikebukuro T, Mizoguchi H, Takakura N. Effects of acupuncture and heating on blood volume and oxygen saturation of human Achilles tendon *in vivo*. *Eur J Appl Physiol.* 2010;**109**(3):545–50.
- 166 Larsson R, Oberg PA, Larsson SE. Changes of trapezius muscle blood flow and electromyography in chronic neck pain due to trapezius myalgia. *Pain.* 1999;**79**(1):45–50.
- 167 Sandberg M, Lundeberg T, Lindberg LG, Gerdle B. Effects of acupuncture on skin and muscle blood flow in healthy subjects. *Eur J Appl Physiol.* 2003;**90**(1–2):114–9.
- 168 Shinbara H, Okubo M, Sumiya E, Fukuda F, Yano T, Kitade T. Effects of manual acupuncture with sparrow pecking on muscle blood flow of normal and denervated hindlimb in rats. *Acupunct Med.* 2008;**26**(3):149–59.
- 169 Larsson R, Cai H, Zhang Q, Oberg PA, Larsson SE. Visualization of chronic neck-shoulder pain: impaired microcirculation in the upper trapezius muscle in chronic cervicobrachial pain. *Occup Med (Lond).* 1998;**48**(3):189–94.
- 170 James SL, Ali K, Pocock C, Robertson C, Walter J, Bell J, et al. Ultrasound guided dry needling and autologous blood

- injection for patellar tendinosis. *Br J Sports Med.* Aug 2007;**41**(8):518–21; discussion 522.
- 171 Lee JA, Jeong HJ, Park HJ, Jeon S, Hong SU. Acupuncture accelerates wound healing in burn-injured mice. *Burns.* 2011;**37**(1):117–25.
 - 172 Pfeifer MT, Cooper SR, Uhl NL. Chiropractic management of tendinopathy: a literature synthesis. *J Manipulative Physiol Ther.* 2009;**32**(1):41–52.
 - 173 Cook J, Purdham C. Is tendon pathology a continuum? A pathology based model to explain the clinical presentation of load induced tendinopathy. *Br J Sports Med.* 2009;**43**(6):409–16.
 - 174 Kietrys DM, Palombaro KM, Azzaretto E, Hubler R, Schaller B, Schluskel JM, *et al.* Effectiveness of dry needling for upper quarter myofascial pain: a systematic review and meta-analysis. *J Orthop Sports Phys Ther.* 2013;**43**(9):620–634.
 - 175 Fernandez-Carnero J, Fernandez-de-las-Penas C, Sterling M, Souvlis T, Arendt-Nielsen L, Vicenzino B. Exploration of the extent of somato-sensory impairment in patients with unilateral lateral epicondylalgia. *J Pain.* 2009;**10**(11):1179–85.
 - 176 Gonzalez-Iglesias J, Cleland JA, del Rosario Gutierrez-Vega M, Fernandez-de-las-Penas C. Multimodal management of lateral epicondylalgia in rock climbers: a prospective case series. *J Manipulative Physiol Ther.* 2011;**34**(9):635–42.
 - 177 Wainner RS, Whitman JM, Cleland JA, Flynn TW. Regional interdependence: a musculoskeletal examination model whose time has come. *J Orthop Sports Phys Ther.* Nov 2007;**37**(11):658–660.
 - 178 Bergman GJ, Winters JC, Groenier KH, *et al.* Manipulative therapy in addition to usual medical care for patients with shoulder dysfunction and pain: a randomized, controlled trial. *Ann Intern Med.* Sep 21 2004;**141**(6):432–439.
 - 179 Mintken PE, Cleland JA, Carpenter KJ, Bieniek ML, Keirns M, Whitman JM. Some factors predict successful short-term outcomes in individuals with shoulder pain receiving cervicothoracic manipulation: a single-arm trial. *Phys Ther.* Jan 2010;**90**(1):26–42.
 - 180 Sobel JS, Kremer I, Winters JC, Arendzen JH, de Jong BM. The influence of the mobility in the cervicothoracic spine and the upper ribs (shoulder girdle) on the mobility of the scapulohumeral joint. *J Manipulative Physiol Ther.* Sep 1996;**19**(7):469–474.
 - 181 Sobel JS, Winters JC, Groenier K, Arendzen JH, Meyboom de Jong B. Physical examination of the cervical spine and shoulder girdle in patients with shoulder complaints. *J Manipulative Physiol Ther.* May 1997;**20**(4):257–262.
 - 182 Strunce JB, Walker MJ, Boyles RE, Young BA. The immediate effects of thoracic spine and rib manipulation on subjects with primary complaints of shoulder pain. *J Man Manip Ther.* 2009;**17**(4):230–236.
 - 183 Winters JC, Sobel JS, Groenier KH, Arendzen HJ, Meyboom-de Jong B. Comparison of physiotherapy, manipulation, and corticosteroid injection for treating shoulder complaints in general practice: randomised, single blind study. *Bmj.* May 3 1997;**314**(7090):1320–1325.
 - 184 Bang MD, Deyle GD. Comparison of supervised exercise with and without manual physical therapy for patients with shoulder impingement syndrome. *J Orthop Sports Phys Ther.* Mar 2000;**30**(3):126–137.
 - 185 Boyles RE, Ritland BM, Miracle BM, *et al.* The short-term effects of thoracic spine thrust manipulation on patients with shoulder impingement syndrome. *Man Ther.* Aug 2009;**14**(4):375–380.
 - 186 Cotchett MP, Landorf KB, Munteanu SE, Raspovic AM. Consensus for dry needling for plantar heel pain (plantar fasciitis): a modified Delphi study. *Acupunct Med.* 2011;**29**(3):193–202.
 - 187 Cotchett MP, Landorf KB, Munteanu SE, Raspovic A. Effectiveness of trigger point dry needling for plantar heel pain: study protocol for a randomised controlled trial. *J Foot Ankle Res.* 2011;**4**:5.
 - 188 Muller R, Giles LG. Long-term follow-up of a randomized clinical trial assessing the efficacy of medication, acupuncture, and spinal manipulation for chronic mechanical spinal pain syndromes. *J Manipulative Physiol Ther.* 2005;**28**(1):3–11.
 - 189 He D, Veiersted KB, Hostmark AT, Medbo JI. Effect of acupuncture treatment on chronic neck and shoulder pain in sedentary female workers: a 6-month and 3-year follow-up study. *Pain.* 2004;**109**(3):299–307.
 - 190 Fernandez-Carnero J, Cleland JA, Arbizu RL. Examination of motor and hypoalgesic effects of cervical vs thoracic spine manipulation in patients with lateral epicondylalgia: a clinical trial. *J Manipulative Physiol Ther.* 2011;**34**(7):432–40.
 - 191 Struijs PA, Damen PJ, Bakker EW, Blankevoort L, Assendelft WJ, van Dijk CN. Manipulation of the wrist for management of lateral epicondylitis: a randomized pilot study. *Phys Ther.* 2003;**83**(7):608–16.
 - 192 Brantingham JW, Lee Gilbert J, Shaik J, Globe G. Sagittal plane blockage of the foot, ankle and hallux and foot alignment-prevalence and association with low back pain. *J Chiropr Med.* 2006;**5**(4):123–7.
 - 193 Cambron JA, Duarte M, Dexheimer J, Solecki T. Shoe orthotics for the treatment of chronic low back pain: a randomized controlled pilot study. *J Manipulative Physiol Ther.* 2011;**34**(4):254–60.
 - 194 Cibulka MT. Low back pain and its relation to the hip and foot. *J Orthop Sports Phys Ther.* 1999;**29**(10):595–601.
 - 195 Dananberg HJ, Guiliano M. Chronic low-back pain and its response to custom-made foot orthoses. *J Am Podiatr Med Assoc.* 1999;**89**(3):109–17.
 - 196 Williams AE, Hill LA, Nester CJ. Foot orthoses for the management of low back pain: a qualitative approach capturing the patient's perspective. *J Foot Ankle Res.* 2013;**6**(1):17.